

Dental anomalies in an orthodontic patient population with maxillary lateral incisor agenesis

Mehmet Citak¹, Elif Bahar Cakici², Yasin Atakan Benkli³, Fatih Cakici², Bircan Bektas⁴, Suleyman Kutalmış Buyuk³

DOI: <http://dx.doi.org/10.1590/2177-6709.21.6.098-102.oar>

Introduction: The purpose of this study was to evaluate the prevalence of dental anomalies in a subpopulation of orthodontic patients with agenesis of maxillary lateral incisors (MLI).

Methods: The material of the present study included the records of the 1964 orthodontic patients. Panoramic radiographs and dental casts were used to analyze other associated eight dental anomalies, including agenesis of other teeth, *dens invaginatus*, *dens evaginatus*, peg shaped MLI, taurodontism, pulp stone, root dilaceration and maxillary canine impaction.

Results: Out of the 1964 patients examined, 90 were found to have agenesis of MLI, representing a prevalence of 4.6%. The most commonly found associated-anomalies were agenesis of other teeth (23.3%), peg-shaped MLIs (15.6%), taurodontism (42.2%), and dilacerated teeth (18.9%).

Conclusion: Permanent tooth agenesis, taurodontism, peg-shaped maxillary lateral incisor, and root dilacerations are frequently associated with maxillary lateral incisor agenesis.

Keywords: Dental anomalies. Hypodontia. Panoramic radiograph.

¹Research Assistant, Department of Endodontics, Faculty of Dentistry, Ordu University, Ordu, Turkey.

²Assistant Professor, Department of Endodontics, Faculty of Dentistry, Ordu University, Ordu, Turkey.

³Assistant Professor, Department of Orthodontics, Faculty of Dentistry, Ordu University, Ordu, Turkey.

⁴Research Assistant, Department of Orthodontics, Faculty of Dentistry, Ordu University, Ordu, Turkey.

How to cite this article: Citak M, Cakici EB, Benkli YA, Cakici F, Bektas B, Buyuk SK. Dental anomalies in an orthodontic patient population with maxillary lateral incisor agenesis. Dental Press J Orthod. 2016 Nov-Dec;21(6):98-102. DOI: <http://dx.doi.org/10.1590/2177-6709.21.6.098-102.oar>

Submitted: March 18, 2016 - **Revised and accepted:** July 20, 2016

» The authors report no commercial, proprietary or financial interest in the products or companies described in this article.

» Patients displayed in this article previously approved the use of their facial and intraoral photographs.

Contact address: Suleyman Kutalmış Buyuk
E-mail: skbuyuk@gmail.com

INTRODUCTION

Dental anomalies are typically caused by either genetic or environmental stimuli.^{1,2} Mutations in AXIN2, PAX9 and MSX1 have been determined in families with dental agenesis.^{3,4} The most frequently observed dental agenesis in children is defined as the absence of one or more primary/permanent teeth.⁵ Data for congenital tooth agenesis prevalence vary between 0.3 and 11.3%^{6,7} for both males and females. However, the prevalence of congenital tooth agenesis was shown to be higher in females than in males, in some reports.^{5,8,9}

After third molars, maxillary lateral incisors (MLI) are the teeth that are the most frequently missing.^{5,10} Agenesis of MLI has been documented for its higher prevalence than of other permanent teeth.¹¹ A correlation between MLI agenesis and palatally displaced canines,¹² tooth transpositions,¹³ and premolar rotations¹⁴ has also been reported. However, reports on the prevalence of dental anomalies in a large MLI agenesis patient cohort have not been determined. The purpose of the current study was to investigate the prevalence of MLI agenesis and other dental anomalies in an orthodontic subpopulation in Turkey.

MATERIAL AND METHODS

Panoramic radiographs of 1964 patients (1174 females, 790 males) of the Department of Orthodontics of Ordu University, Turkey, between January 2013 and September 2015, were retrospectively analyzed. Patients, aged 12 to 25 years, with unilateral or bilateral agenesis of MLI and panoramic radiograph were included in the study. Patients with incomplete records, permanent tooth extraction, and/or poor-quality

panoramic radiographs were excluded from the study. Care was used to ensure that all radiographs were taken by the same technician operating the same panoramic roentgen unit device (Kodak Cephalostat, Rochester NY, USA). In order to eliminate inter-examiner differences, all records were examined by one observer. All radiographs were evaluated by an orthodontist with more than 15 years of experience.

The following anomalies were determined in this study (Fig 1):

- 1) Agenesis: congenital developmental loss of one or more permanent teeth.
- 2) *Dens invaginatus*: caused by the invagination of enamel into the dental papilla before the mineralization phase.¹⁵
- 3) *Dens evaginatus*: malformation characterized by an accessory cusp, composed of normal enamel and dentine, with or without pulp tissue.¹⁶
- 4) Microdontia (peg-shaped teeth): teeth that are substantially smaller than the average normal size. Microdontia also refers to a tooth that does not fill its space in the dental arch or appears small due to absence of expected shape.¹⁷
- 5) Taurodontism: vertically extended, extremely oversized pulp cavities that are apically displaced at the pulpal floor.¹⁸
- 6) Pulp stone: calcified masses on the pulp of healthy, diseased, and even unerupted teeth freely attached or embedded into the coronal rather than the root portion of the pulp organ.¹⁹
- 7) Dilaceration: deviation in the linear relationship of a crown of a tooth to its root.²⁰




Figure 1 - A) Root dilacerations. B) Pulp stone. C) Taurodontism. D) *Dens evaginatus*. E) *Dens invaginatus*.

8) Impaction: a tooth that is predicted to remain unerupted because of a physical barrier or deflection along its eruption path.²¹

Statistical analysis

Statistics were calculated with SPSS 15.0 statistical software (SPSS Inc, Chicago, IL, USA). Anomaly prevalence was measured with respect to sex, side and dental location. Chi-square analysis and MLI prevalence were compared to previously published reports from the Turkish population.²²⁻²⁸ A p-value of less than 0.05 was considered significant.

To examine errors associated with digitizing and measurements, 10% of panoramic images were selected randomly and all dental anomalies were evaluated by the same author four weeks after the first examination. Kappa coefficients were used to calculate the reliability of each dental anomaly determination from the two evaluation periods.

RESULTS

Kappa score of each dental anomaly was 1.00. This score indicated good agreement with the first and second

evaluations and was observed for each dental anomaly. Out of the 1964 subjects (1174 females, 790 males) evaluated, 90 (62 females, 28 males) were determined to have MLI agenesis (prevalence = 4.58%; being 5.3% for females and 3.5% males. Difference between males and females was statistically not significant [$X^2 = 3.26$; $p = 0.071$]). Bilateral MLI agenesis was found in 62 subjects (68.9%) and unilateral agenesis in 28 patients (31.1%).

The investigated dental anomalies in MLI agenesis patients were: *dens invaginatus*, *dens evaginatus*, peg-shaped MLI, taurodontism, pulp stone, root dilacerations, impaction of maxillary canine, and missing teeth other than third molars (Table 1). The prevalence of MLI agenesis-associated dental-anomalies was referenced to previous work, for consistency (Table 2). The prevalence of agenesis of other teeth ($p < 0.001$), peg-shaped MLIs ($p < 0.001$), taurodontism ($p < 0.001$), and dilacerated teeth ($p < 0.01$) were greater in our sample compared to the other studies. No statistical significant difference was shown for the prevalence of *dens invaginatus* ($p = 0.888$), *dens evaginatus* ($p = 0.123$), pulp stone ($p = 0.666$) and impaction of maxillary canines ($p = 0.477$).

Table 1 - Distribution of MLI agenesis-associated dental-anomalies.

Dental anomaly	Male / Female	Unilateral / Bilateral	Total (%)
Agenesis of other teeth	5/16	7/14	21 (23.3)
<i>Dens invaginatus</i>	1/0	1/0	1 (1.1)
<i>Dens evaginatus</i>	1/1	1/1	2 (2.2)
Peg shaped MLI	7/7	14/0	14 (15.6)
Taurodontism	9/29	1/37	38 (42.2)
Pulp stone	0/9	5/4	9 (10)
Dilaceration	4/13	5/12	17 (18.9)
Impaction of maxillary canine	2/3	3/2	5 (5.6)

Table 2 - Comparison of the frequencies of dental anomalies subjects with maxillary lateral incisor agenesis and previous studies.

Dental anomaly	Present study			Reference studies			p*
	n	%	n	Total	%	Literature	
Agenesis of other teeth	21	23.3	144	3165	5.0	Kazanci et al. ²²	< 0.001
<i>Dens invaginatus</i>	1	1.1	13	1012	1.3	Cakici et al. ²³	0.888
<i>Dens evaginatus</i>	2	2.2	56	900	6.2	Uslu et al. ²⁴	0.123
Peg shaped MLI	14	15.6	46	3043	1.5	Altug-Atac, Erdem ²⁵	< 0.001
Taurodontism	38	42.2	9	900	1.0	Uslu et al. ²⁴	< 0.001
Pulp stone	9	10	60	519	11.6	Gulsahi et al. ²⁶	0.666
Dilaceration	17	18.9	214	2251	9.5	Miloglu et al. ²⁷	0.003
Impaction of maxillary canine	5	5.6	488	12000	4.1	Gunduz, Celenk ²⁸	0.477

n = number of subjects, MLI; Maxillary lateral incisor, * p indicates results of chi-square test.

DISCUSSION

The prevalence of dental anomalies is variable among different populations. The aim of the present study was to determine the prevalence of MLI agenesis-associated dental anomalies in orthodontic Turkish patients. We found that about 4.58% of patients had one or both maxillary incisors missing. These results are consistent with that reported by others, which ranges from 0.3% and 11.3%.^{9,10,25} The prevalence of MLI agenesis varies considerably between studies.^{7,29-32} Horowitz³³ showed a prevalence of 1.11% in an adolescent population (n=1000; ages ranging from 7 to 16 years). Celikoglu et al⁷ reported a prevalence of 2.4% from 3872 East Anatolian adolescent patients in Turkey. The differences may be related to sample size or selection, but may be different due to regional ethnic population, genetic variability, and/or environmental factors.

There was no statistically significant correlation between sex and MLI agenesis. Interestingly, we had more female subjects in our cohort. Some reports have shown insignificant differences,^{34,35} while others have determined significant sex-related changes.^{36,37}

Previous studies^{24,38-41} have reported that tooth agenesis can be associated with other dental malformations, such as taurodontism, transposition, microdontia, ectopic eruption, supernumerary tooth or peg-shaped MLI. Most of the papers^{8,42,43} published on MLI agenesis demonstrated a reduction in crown size or a peg-shaped form of the contralateral MLI. MLI agenesis was detected more commonly in females than males.

When we compared the prevalence rates of MLI agenesis-associated dental anomalies and reference values,²²⁻²⁸ it was determined that the prevalence rates were significantly augmented for taurodontism, agenesis of other teeth, and peg-shaped MLIs. There have been only two studies that have compared their work with^{7,38} reference values. However, taurodontism, pulp stone, *dens invaginatus*, *dens evaginatus*, and impaction of the maxillary canine were not assessed in those studies. In this respect, our study is the first one to show the different associated dental anomalies between subjects with MLI agenesis.

The most common MLI agenesis-associated dental anomaly was taurodontism; in our study, with a prevalence of 42.2%. Uslu et al²⁴ showed 1% of taurodontism prevalence in 900 orthodontic patients. Discrepancy in the results may be related to the location variations, and our taurodontism detection method. The difference

might arise from racial differences or differences in diagnostic criteria.

Out of the 28 patients who had unilateral absence, 14 (50.0%) were found to have a peg-shaped lateral incisor on the other side. Altug-Atac and Erdem²⁵ reported that 1.51% of patients had peg-shaped MLI in an orthodontic patient population. However, Albashaireh and Khaider⁴² demonstrated that peg-shaped and reduced size maxillary lateral incisors were found in 2.3% and 2.9% of patients, respectively. Similar to the prevalence of our results, Albashaireh and Khaider⁴² showed 50% microdontia or peg-shaped MLIs on the other side in individuals with unilateral MLI agenesis. In this study, the prevalence of other teeth agenesis (23.3%) was very high, compared to the reports by Celikoglu et al.⁷ The higher rate reported may be attributed to orthodontic malocclusions, which supports the findings by Garib et al.³⁸

The prevalence of root dilacerations in our study (18.9%) was greater than that reported of a reference study from the general population (9.5%).²⁷ Diagnosing dilacerations is mostly imperative for root canal treatment, tooth extraction, and Orthodontics.³⁴ The other associated anomalies (*dens invaginatus*, *dens evaginatus*, pulp stone and impaction of maxillary canine prevalence) were similarly found with reference studies.^{23,24,26,28}

Associations between tooth anomalies are clinically relevant, and early diagnosis may be helpful to reduce risk.⁴⁴ Therefore, diagnosis and treatment options should be precisely made. We found a higher prevalence of associated dental anomalies in MLI agenesis patients in Turkish orthodontic population. However, the orthodontic literature shows different prevalence rates of dental anomalies from the general population.^{7,9,22,25,34} This is mostly likely due to the greater variability of racial factors, environmental stimuli and genetics.

CONCLUSIONS

The increased prevalence of MLI agenesis-associated dental anomalies was validated by previous reports from another orthodontic patient population. There was a significant correlation between MLI agenesis and the agenesis of other permanent teeth. In addition, increased agenesis of other teeth, taurodontism, peg-shaped maxillary lateral incisor and root dilacerations were also statistically significant. These associations can be most likely explained by genetic or environmental factors that may contribute to these dental anomalies.

Author contributions

Conception/design of the study: SKB, YAB, FC.
Acquisition, analysis or interpretation: YAB, BB,

MC, SKB. Wrote the article: SKB, EBC. Critical revision of the article: SKB, YAB. Final approval of the article: SKB, EBC.

REFERENCES

- Akcam MO, Evirgen S, Uslu O, Memikoğlu UT. Dental anomalies in individuals with cleft lip and/or palate. *Eur J Orthod*. 2010 Apr;32(2):207-13.
- Kotsomitis N, Dunne MP, Freer TJ. A genetic aetiology for some common dental anomalies: a pilot twin study. *Aust Orthod J*. 1996 Oct;14(3):172-8.
- Nieminen P. Genetic basis of tooth agenesis. *J Exp Zool B Mol Dev Evol*. 2009 Jun 15;312B(4):320-42.
- De Coster PJ, Marks LA, Martens LC, Huysseune A. Dental agenesis: genetic and clinical perspectives. *J Oral Pathol Med*. 2009 Jan;38(1):1-17.
- Celikoglu M, Kazanci F, Miloglu O, Oztek O, Kamak H, Ceylan I. Frequency and characteristics of tooth agenesis among an orthodontic patient population. *Med Oral Patol Oral Cir Bucal*. 2010 Sept 1;15(5):e797-801.
- O'Dowling IB, McNamara TG. Congenital absence of permanent teeth among Irish school-children. *J Ir Dent Assoc*. 1990;36(4):136-8.
- Celikoglu M, Kamak H, Yildirim H, Ceylan I. Investigation of the maxillary lateral incisor agenesis and associated dental anomalies in an orthodontic patient population. *Med Oral Patol Oral Cir Bucal*. 2012 Nov 1;17(6):e1068-73.
- Pinho T, Tavares P, Maciel P, Pollmann C. Developmental absence of maxillary lateral incisors in the Portuguese population. *Eur J Orthod*. 2005 Oct;27(5):443-9. Epub 2005 Aug 31.
- Silva Meza R. Radiographic assessment of congenitally missing teeth in orthodontic patients. *Int J Paediatr Dent*. 2003 Mar;13(2):112-6.
- Fekonja A. Hypodontia in orthodontically treated children. *Eur J Orthod*. 2005 Oct;27(5):457-60.
- Polder BJ, Van't Hof MA, Van der Linden FP, Kuijpers-Jagtman AM. A meta-analysis of the prevalence of dental agenesis of permanent teeth. *Community Dent Oral Epidemiol*. 2004 June;32(3):217-26.
- Sacerdoti R, Baccetti T. Dentoskeletal features associated with unilateral or bilateral palatal displacement of maxillary canines. *Angle Orthod*. 2004 Dec;74(6):725-32.
- Peck S, Peck L, Kataja M. Concomitant occurrence of canine malposition and tooth agenesis: evidence of orofacial genetic fields. *Am J Orthod Dentofacial Orthop*. 2002 Dec;122(6):657-60.
- Baccetti T. Tooth rotation associated with aplasia of nonadjacent teeth. *Angle Orthod*. 1998 Oct;68(5):471-4.
- Oehlers FA. Dens invaginatus (dilated composite odontome). I. Variations of the invagination process and associated anterior crown forms. *Oral Surg Oral Med Oral Pathol*. 1957 Nov;10(11):1204-18.
- Levitani ME, Himel VT. Dens evaginatus: literature review, pathophysiology, and comprehensive treatment regimen. *J Endod*. 2006 Jan;32(1):1-9.
- Kocabalkan E, Ozyemişci N. Restoration of severe hypodontia associated with microdontia by using an overdenture: a clinical report. *Chin Med J (Engl)*. 2005 Feb 20;118(4):350-2.
- Shifman A, Channanell I. Prevalence of taurodontism found in radiographic dental examination of 1,200 young adult Israeli patients. *Community Dent Oral Epidemiol*. 1978 July;6(4):200-3.
- Al-Hadi Hamasha A, Darwazeh A. Prevalence of pulp stones in Jordanian adults. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1998 Dec;86(6):730-2.
- Hamasha AA, Al-Khateeb T, Darwazeh A. Prevalence of dilaceration in Jordanian adults. *Int Endod J*. 2002 Nov;35(11):910-2.
- Thilander B, Jakobsson SO. Local factors in impaction of maxillary canines. *Acta Odontol Scand*. 1968 May;26(2):145-68.
- Kazanci F, Celikoglu M, Miloglu O, Ceylan I, Kamak H. Frequency and distribution of developmental anomalies in the permanent teeth of a Turkish orthodontic patient population. *J Dent Sci*. 2011 June;6(2):82-9.
- Cakici F, Celikoglu M, Arslan H, Topcuoglu HS, Erdogan AS. Assessment of the prevalence and characteristics of dens invaginatus in a sample of Turkish Anatolian population. *Med Oral Patol Oral Cir Bucal*. 2010 Nov 1;15(6):e855-8.
- Uslu O, Akcam MO, Evirgen S, Cebeci I. Prevalence of dental anomalies in various malocclusions. *Am J Orthod Dentofacial Orthop*. 2009 Mar;135(3):328-35.
- Altug-Atac AT, Erdem D. Prevalence and distribution of dental anomalies in orthodontic patients. *Am J Orthod Dentofacial Orthop*. 2007 Apr;131(4):510-4.
- Gulsahi A, Cebeci AI, Ozden S. A radiographic assessment of the prevalence of pulp stones in a group of Turkish dental patients. *Int Endod J*. 2009 Aug;42(8):735-9.
- Miloglu O, Cakici F, Caglayan F, Yilmaz AB, Demirkaya F. The prevalence of root dilacerations in a Turkish population. *Med Oral Patol Oral Cir Bucal*. 2010 May 1;15(3):e441-4.
- Gunduz K, Celenk P. The incidence of impacted transmigrant canines: a retrospective study. *Oral Radiol*. 2010 Dec;26(2):77-81.
- Baccetti T. A controlled study of associated dental anomalies. *Angle Orthod*. 1998 June;68(3):267-74.
- Johannsdottir B, Wisth PJ, Magnusson TE. Prevalence of malocclusion in 6-year-old Icelandic children. *Acta Odontol Scand*. 1997 Dec;55(6):398-402.
- Muller TP, Hill IN, Peterson AC, Blayney JR. A survey of congenitally missing permanent teeth. *J Am Dent Assoc*. 1970 July;81(1):101-7.
- Rolling S. Hypodontia of permanent teeth in Danish schoolchildren. *Scand J Dent Res*. 1980 Oct;88(5):365-9.
- Horowitz JM. Aplasia and malocclusion: a survey and appraisal. *Am J Orthod*. 1966 June;52(6):440-53.
- Thongudomporn U, Freer TJ. Prevalence of dental anomalies in orthodontic patients. *Aust Dent J*. 1998 Dec;43(6):395-8.
- Endo T, Ozoe R, Kubota M, Akiyama M, Shimooka S. A survey of hypodontia in Japanese orthodontic patients. *Am J Orthod Dentofacial Orthop*. 2006 Jan;129(1):29-35.
- Basdra EK, Kiokpasoglu M, Stellzig A. The Class II Division 2 craniofacial type is associated with numerous congenital tooth anomalies. *Eur J Orthod*. 2000 Oct;22(5):529-35.
- Bergström K. An orthopantomographic study of hypodontia, supernumeraries and other anomalies in school children between the ages of 8-9 years. An epidemiological study. *Swed Dent*. 1977;1(4):145-57.
- Garib DG, Alencar BM, Lauris JR, Baccetti T. Agenesis of maxillary lateral incisors and associated dental anomalies. *Am J Orthod Dentofacial Orthop*. 2010 June;137(6):732.e1-6.
- Gomes RR, da Fonseca JA, Paula LM, Faber J, Acevedo AC. Prevalence of hypodontia in orthodontic patients in Brasília, Brazil. *Eur J Orthod*. 2010 June;32(3):302-6.
- Shapira Y, Kuftinec MM. Maxillary tooth transpositions: characteristic features and accompanying dental anomalies. *Am J Orthod Dentofacial Orthop*. 2001 Feb;119(2):127-34.
- Zhu JF, Marcushamer M, King DL, Henry RJ. Supernumerary and congenitally absent teeth: a literature review. *J Clin Pediatr Dent*. 1996 Winter;20(2):87-95.
- Albashaieh ZS, Khader YS. The prevalence and pattern of hypodontia of the permanent teeth and crown size and shape deformity affecting upper lateral incisors in a sample of Jordanian dental patients. *Community Dent Health*. 2006 Dec;23(4):239-43.
- Woolf CM. Missing maxillary lateral incisors: a genetic study. *Am J Hum Genet*. 1971 May;23(3):289-96.
- Apajalahti S, Holtta P, Turtola L, Pirinen S. Prevalence of short-root anomaly in healthy young adults. *Acta Odontol Scand*. 2002 Jan;60(1):56-9.